



Cutoff low and landfalling AR bring heavy rainfall to the southwestern U.S.

- A quasi-stationary cutoff low interacted with a region of tropical moisture over the Eastern Pacific Ocean, resulting in a long-duration atmospheric river (AR) over the Baja Peninsula
- Poleward moisture transport and strong synoptic-dynamic forcing for ascent produced multiple episodes of heavy rainfall across Southern California, Arizona, and southern Nevada

Elevated sections of Southern California and central Arizona received 3–6 inches of total precipitation, with 1–3 inches at lower


elevations








Coastal AR Scale Analysis







- A landfalling AR brought AR3/AR4 conditions to the Baja Peninsula, with some places experiencing AR conditions for more than 72 hours
- The total time-integrated IVT during this event was a remarkable 13.76 × 10⁷ kg m⁻¹ at 27°N, 114°W
- The rapid increase in IVT values after 1800 UTC 11 Mar preceded the 24-hour period (1200 UTC 12–13 Mar) with the most intense rainfall in Southern California and western Arizona



Inland AR Scale Analysis






Prolonged inland penetration of high IVT values resulted in AR2 conditions over southern Arizona, with some areas experiencing AR conditions for more than 72 hours




GFS Analyses: Valid 0000 UTC 11 Mar



- This AR formed as a result of the interaction between a cutoff low west of Southern California and a region of deep tropical moisture over the Eastern Pacific Ocean
- 0000 UTC 11 Mar GFS analysis indicates that the first episode of heavy rainfall was associated with strong low-to-midlevel southwesterly flow downstream of the cutoff low that transported very moist air over extreme southeastern California and southwestern Arizona




GFS Analyses: Valid 0000 UTC 13 Mar



- As time progressed, the cutoff low slowly propagated eastward, and the moisture plume began to dissipate
- 0000 UTC 13 Mar GFS analysis suggests that the second episode of heavy rainfall was associated with a secondary region of enhanced and cyclonically curved IVT that developed immediately east of the cutoff low
- During both episodes of heavy rainfall, synoptic-dynamic forcing for ascent (via differential cyclonic vorticity advection) also played an important role







- The AR associated with the cutoff low transported deep moisture into the Desert Southwest, with precipitable water (PWAT) values exceeding 4 standard deviations above normal across Arizona and New Mexico
- The 1200 UTC 11 Mar atmospheric sounding from Tucson, AZ (TUS), shows nearly saturated conditions between 800-hPa and 550-hPa
- TUS set a new record March PWAT value of 1.14 inches (> 300% of normal)



CNRFC 120-h Observed (Raw) Precipitation: Valid 9 AM PDT 14 Mar






Source: NOAA/NWS CNRFC, https://www.cnrfc.noaa.gov/

Source: NOAA/NWS Advanced Hydrologic Prediction Service, https://water.weather.gov/ahps/

- The heaviest precipitation (3–6 inches) fell across the Transverse and Peninsular Ranges in Southern California, as well as the higher terrain in west-central Arizona
- Coastal, valley, and desert areas received about 1-3 inches of total rainfall
- McCarran International Airport (KLAS) and Yuma International Airport (KYUM) both set daily precipitation records on 10 Mar and 12 Mar
- North Las Vegas Airport (KVGT) and KYUM recorded more than 50% of their average annual rainfall from this event






Source: NOAA/NWS Colorado Basin River Forecast Center, https://www.cbrfc.noaa.gov/

- NWS Phoenix received numerous reports of flash flooding in Imperial County, CA, and Maricopa County, AZ, on 12–13 Mar
- Heavy rainfall triggered a rapid streamflow response on many rivers in western and central Arizona during the morning of 13 Mar
- The Agua Fria River (near Rock Springs, AZ) rose more than 10 feet in a 6-hour period and briefly exceeded moderate flood stage






Source: NOAA/NWS Colorado Basin River Forecast Center, https://www.cbrfc.noaa.gov/

• The Big Sandy River (near Wikieup, AZ) rose more than 11 feet in a 3-hour period and recorded its all-time maximum stage height (19.06 ft) and its highest peak discharge (33,400 cfs) over the past 10 water years (since 1 Oct 2010)






Source: NOAA/NWS Colorado Basin River Forecast Center, https://www.cbrfc.noaa.gov/

• The Verde River (near Clarkdale, AZ) recorded its second-highest peak discharge (10,700 cfs) and daily mean discharge (5,550 cfs) over the past 10 water years





Source: NOAA/NWS Colorado Basin River Forecast Center, https://www.cbrfc.noaa.gov/

• The Salt River (near Roosevelt, AZ) recorded its highest peak discharge (20,300 cfs) and daily mean discharge (14,400 cfs) over the past 10 water years